

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** AFCJB33010

**Data Sensitivity:** No

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Plagopterus argentissimus*

**COMMON NAME:** Woundfin

**SYNONYMS:** *Meda argentissimus*

**FAMILY:** Cyprinidae

**AUTHOR, PLACE OF PUBLICATION:** Cope, 1874. Proc. Am. Philos. Soc. 14:129-39.

**TYPE LOCALITY:** San Luis Valley, western Colorado [sic]; Virgin River, Washington County, southwestern Utah (Miller and Hubbs, 1960).

**TYPE SPECIMEN:**

**TAXONOMIC UNIQUENESS:** Monotypic genus. Tribe of 6 species endemic to the Lower Colorado River basin. As referenced in Minckley (1973): Miller and Hubbs (1960) reviewed the tribe Plagopterini and concluded that *Plagopterus* is the most advanced genus among the three represented, and that it and *Meda* are the most closely related. Uyeno and Miller (1973) reached the same conclusions based upon karyotypes, as did Coad (1976 in Lee et al. 1980) based on a study of 28 characteristics. No described subspecies.

**DESCRIPTION:** From Minckley (1973): "Rarely exceeds 75.0 mm (2.95 in.)....Body thicker anteriorly, thinner posteriorly, flattened beneath. Scales represented by dermal plates supporting low ridges on antero-dorsal portion of body, deeply embedded. First spinous ray strong, sharp-pointed. Barbels present. Dorsal fin-rays 8 or 9. Anal fin-rays usually 10. Pharyngeal teeth typically 1, 5-4, 1.... Coloration silvery over-all; sometimes with watery-yellow at bases of pectoral fins, and developing ventro-lateral, pinkish coloration in exceedingly "high," reproductive males."

**AIDS TO IDENTIFICATION:** A small slender, silvery, scaleless minnow. Head and belly flattened, and mouth small and nearly horizontal. As with the Virgin spinedace and the spikedace, the woundfin has two large spines in dorsal fin and bright silver sides. No scales. Long snout; barbel at mouth corner. Can be distinguished from spikedace and spinedace by the presence of barbels. Woundfin has wider, flatter head than spikedace and lacks the scales seen in spinedace. In its present range, the woundfin is unlikely to be confused with sympatric species.

**ILLUSTRATIONS:** B&W photo (Minckley 1973:115)

Color drawing (Page and Burr 1991)

**TOTAL RANGE:** Historic range includes the lower Colorado River basin including the Virgin, Moapa, Salt and Gila river systems. At present, the woundfin is totally restricted to approximately 50 miles of perennial reaches of the Virgin River in the states of Utah, Arizona, and Nevada.

**RANGE WITHIN ARIZONA:** Found sporadically throughout the Arizona portion of the Virgin River mainstem (Mohave County). Has been documented recently by the Virgin River Fishes Recovery Team near Cedar Pockets in the Virgin River gorge, and near the confluence of the Virgin River and Beaver Dam Wash (both locations are permanent survey sites for the team's biannual monitoring program).

### **SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** This species is able to exist in heavily mineralized waters and can tolerate extremely high turbidity. Found in association with Virgin chub, *Gila seminuda*. Appears to undertake relatively long downstream migrations within present habitat.

**REPRODUCTION:** Fry generally appear by June, but continue to be produced throughout summer. Maturation appears to occur in second summer, and few if any individuals live more than four years. Has been successfully spawned at Dexter National Fish Hatchery and Technology Center in New Mexico.

**FOOD HABITS:** Omnivorous; eats algae, detritus, seeds and aquatic insects.

**HABITAT:** The species lives in swift parts of silty streams, seemingly avoiding clear waters and very seldom found in quieter pools. Occupies main channel of seasonally swift, highly turbid, and extremely warm streams, with sandy, constantly shifting bottoms. Current velocities and depths preferred by adults are one or two feet per second and eight to 18 inches. Young seek quiet backwaters with sandy substrates.

**ELEVATION:** The Virgin River reaches its lowest elevation in Littlefield, Arizona at approximately 1,900 ft. (580 m) and its highest elevation in the headwaters in Zion National Park, Utah at approximately 10,000 ft. (3,050 m). The woundfin is currently distributed from the lowermost elevations of the Virgin River to approximately 3,000 ft. (915 m) near the confluence with La Verkin Creek.

**PLANT COMMUNITY:** Presently, the Virgin River riparian community consists primarily of *Tamarix* spp. Biotic communities along the Virgin River include the Great Basin and Mohave desert scrub.

**POPULATION TRENDS:** Declining. Historical habitat has been lost due to human impacts including habitat fragmentation, introduction of nonnative species and dewatering due to

agriculture, mining and urbanization. Damming and drying have caused the disappearance of the woundfin throughout most of its historic range and continue to impact it in the Virgin River. Flows there have been depleted by municipal and agricultural withdrawals. Growth of the town of St. George, Utah, on the Virgin River and in the center of the woundfin's present range will further limit water availability in the area. Additionally, nonnative fishes have been brought to the system, carrying with them parasites previously unknown to woundfin, but now occasionally infesting them. Red shiner, a baitfish, is rapidly expanding its range upstream in the Virgin River and, by still uncertain mechanisms, causing declines in the remaining woundfin populations.

## **SPECIES PROTECTION AND CONSERVATION**

**ENDANGERED SPECIES ACT STATUS:** LE (USDI, FWS 1970)  
Listed Critical Habitat (USDI, FWS 2000)  
Experimental nonessential in portions of the Gila River

**STATE STATUS:** WC (AGFD, WSCA in prep)  
[State Endangered AGFD, TNW 1988]

**OTHER STATUS:** No Forest Service Status (USDA, FS Region 3 1999)  
Forest Service Sensitive (USDA, FS Region 3 1988)  
Endangered, American Fisheries Society.

**MANAGEMENT FACTORS:** Preservation of the last remaining habitat and populations of this species in the Virgin River will require the following: close monitoring of abundance and distribution; guarantees of minimum flows at critical times through critical reaches; control of nonnative fish populations and of parasites introduced with them; refinement of hatchery techniques to produce large numbers for reintroduction; and studies of the species' critical requirements for use in prioritizing management needs in the Virgin River and for evaluating potential reintroduction sites.

**PROTECTIVE MEASURES TAKEN:** Legal statutes that help protect the woundfin include Section 404 of the Clean Water Act and the Endangered Species Act. Virgin River Fishes Recovery Team conducts biannual monitoring and maintains a data base of sampling results. Plans are in process for the construction of fish barriers and eradication of nonnative fish species. Unsuccessful reintroductions have occurred into four historically occupied areas in Arizona (Sycamore Creek on the Prescott N.F., Hassayampa River near Wickenburg, and the Salt and Paria rivers).

**SUGGESTED PROJECTS:** The Virgin River Recovery Team is charged with the mission of restoring the woundfin throughout its former range. The long term goal of the team is the downlisting of the species. Downlisting strategies include the following: restoration and enhancement of existing habitat; establishing and monitoring minimum and channel

maintaining flows; eradication of predatory and competitive nonnative species; reestablishment into and protection of formerly occupied range; critical habitat designation; on-going population monitoring, and other activities as described in the Virgin River Fishes Recovery Plan. Continue to conduct both laboratory and field studies on woundfin genetics, biology, and ecology. Continue to conduct stream surveys to determine the suitability of reintroduction sites.

**LAND MANAGEMENT/OWNERSHIP:** Zion National Park, Bureau of Land Management, U.S. Forest Service, States of Utah, Arizona and Nevada, Washington County Water Conservation District, private, and possibly others.

## **SOURCES OF FURTHER INFORMATION**

### **REFERENCES:**

- Arizona Game and Fish Department. 1988. Threatened Native Wildlife in Arizona. p. 5.
- Arizona Game and Fish Department. In prep. Wildlife of special concern in Arizona.
- Arizona Game and Fish Department Publication. Phoenix, Arizona. 32 pp.
- Cope, E.D. 1874. On the Plagopterinae and the ichthyology of Utah. Amer. Philos. Soc., Proc. 14:129-39.
- Cross, J.N. 1975. Ecological distribution of the fishes of the Virgin River (Utah, Arizona, Nevada) Unpubl. M.S. Thesis, University of Nevada, Las Vegas.
- Deacon, J.E. 1988. The endangered woundfin and water management in the Virgin River, Utah, Arizona, Nevada. Fisheries 13(1):18-24.
- Deacon, J.E. and W.L. Minckley. 1973. A review of information on the woundfin, *Plagopterus argentissimus* Cope (Pisces: Cyprinidae). Progress report on population dispersion and community structure of fishes of the Virgin River System. U.S. Fish and Wildlife Service, Salt Lake City, Utah. pp. 28.
- Hardy, T.B., B. Bartz and W. Carter. 1989. Population dynamics of the fishes in the Virgin River from 1984 through 1987 and impact analyses of the Quail Creek and North Creek reservoir systems. Utah State University. pp. 564.
- Hickman, T.J. 1987. Annotated bibliography for the woundfin, *Plagopterus argentissimus*. Western Ecosystems, P.O. Box 1575, St. George, Utah. pp. 33.
- Miller, R.R. and C.L. Hubbs. 1960. The spiny-rayed cyprinid fishes (Plagopterinei) of the Colorado River system. Misc. Publ. Museum Zoology University Michigan 115:1-39.
- Minckley, W.L. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix. pp. 115-116.
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America, north of Mexico. Houghton Mifflin Co., Boston. p. 83.
- Lee, D.S. et al. 1980. Atlas of North American fishes. North Carolina State Museum of Natural History. pp. 867.
- Sigler, W.F. and R.R. Miller. 1963. Fishes of Utah. Utah State Department of Fish and Game, Salt Lake City.
- USDA, Forest Service Region 3. 1988. Regional Forester's Sensitive Species List.

- USDA, Forest Service Region 3. 1999. Regional Forester's Sensitive Species List.
- USDI, Fish and Wildlife Service. 1970. Federal Register 35:16047.
- USDI, Fish and Wildlife Service 1985. Final rule on proposal to transplant *Plagopterus argentissimus* as an experimental nonessential population in portions of the Gila River Drainage. Federal Register 50:30193.
- USDI, Fish and Wildlife Service. 1992. Virgin River Fishes Recovery Plan. Salt Lake City, Utah. 50 pp.
- USDI, Fish and Wildlife Service. 2000. Endangered and Threatened Wildlife and Plants; Designation of Critical Habitat for the Woundfin and Virgin River Chub. Final Rule. Federal Register 65(17): 4140-4155.
- Uyeno, T. and R. R. Miller. 1973. Chromosomes and the evolution of the Plagoterine fishes (Cyprinidae) of the Colorado River system. Copeia 1973 4:.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

Thomas B. Hardy - Utah State University  
Jim Heinrich - Nevada Division of Wildlife  
Michael Herder - Bureau of Land Management, Arizona Strip  
Leo Lentch - Utah Division of Wildlife Resources  
Henry Maddux - U.S. Fish and Wildlife Service, Salt Lake City

**ADDITIONAL INFORMATION:** As of July 1994, the Arizona Game and Fish Department, Nongame Branch, Native Fish Program is pursuing the reintroduction of the woundfin into the Hassayampa River.

**Revised:** 1994-07-14 (RHB)  
1995-01-29 (KLY)  
2000-01-28 (SMS)

To the user of this abstract: you may use this entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 2001. *Plagopterus argentissimus*. Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.